HOLLY HILL PHARMACY
Home Infusion Services

Holly Hill Pharmacy
1702 Ridgewood Avenue, Suite A - G • Holly Hill, FL 32117
Phone: (386) 677-7377

Penile Injections Diagrammed

Prepare and draw the medication

Use only the dose prescribed for you.


1. Wash your hands.
2. Thaw the medication completely by holding the vial in your hand.
3. Hold the medication bottle so that your fingers do not touch the rubber stopper through which the needle is inserted. Check expiration date of medication.



4. Wipe the top of the vial stopper with an alcohol wipe and let it dry.



5. Remove the needle cover. Do not allow the needle to touch anything prior to drawing the medication or before injecting the medication.



6. Draw an amount of air equal to the amount of medication to be injected into the syringe. Push the needle through the center of the stopper. Push the air into the bottle.



7. Turn the bottle and syringe upside down. Slowly draw the medication into the syringe. Tap the syringe gently to remove the bubbles.



HOLLY HILL PHARMACY
Home Infusion Services

Holly Hill Pharmacy
1702 Ridgewood Avenue, Suite A - G • Holly Hill, FL 32117
Phone: (386) 677-7377


Finish preparation and locate the injection site

8. Move the plunger in and out several times while gently tapping the syringe, thus removing all air bubbles.



9. Gently remove the syringe from the vial.

10. Use the injection site shown by the crosshatch marks in the illustration. Think of the cross-section of the penis as a clock face, with the top at 12 o'clock and the underside at 6 o'clock. Inject at the 10 or 2 o'clock position, not directly on the top or underside.



11. Locate the area of injection. Wipe off with an alcohol swab. Grasp the head of the penis, not the skin. Position the penis along your inner thigh. Maintain traction on the head after cleaning the side of the penis.



HOLLY HILL PHARMACY
Home Infusion Services

Holly Hill Pharmacy
1702 Ridgewood Avenue, Suite A - G • Holly Hill, FL 32117
Phone: (386) 677-7377

Inject and complete the procedure

12. Grasp the syringe between the thumb and middle finger like a pen. Place the needle on the site of injection at a 90-degree angle. Push the needle in, gently but firmly, all the way down to the hub.



13. Shift your finger so that your index finger or your thumb can push in the plunger. Push the medication in slowly - over 8 to 10 seconds.



14. Remove the needle. Apply pressure with your index finger on the injection site and your thumb on the opposite side of the penis. Apply pressure for 2 minutes.



15. Dispose of the used needle and syringe in a sharps container.

16. Return the vial to the freezer.

If you develop priapism

If you have an erection at penetration hardness that lasts for at least 4 hours, seek medical help immediately.